

# Suggestions

1. Separability approximation, Barbanis Hamiltonian and helium excited states.
2. Branch points: their positions (as roots of  $(E_A(\delta) - E_B(\delta))^2$  and a pattern in a complex plane.  
 $1/D^{1/2}$ -expansion for branch points.
3. Prototype Hamiltonian for 2:1 Fermi resonance:  $H = \begin{pmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{pmatrix}$ , where  $H_{11} = \frac{3}{2}\omega_1 + \frac{1}{2}\omega_2$ ,  
 $H_{22} = \frac{1}{2}\omega_1 + \frac{5}{2}\omega_2$ ,  $H_{11} = H_{21} = \frac{\lambda}{2\omega_1^{1/2}\omega_2}$  -- detailed understanding the convergence of linear  
Pade, quadratic, and other algebraic approximants using very high-order calculations.
4. Improving Borel summation by incorporation of the positions of Borel singularities using Borel -  
Darboux approximants  $A(\delta)F(\delta) + B(\delta)(\delta - \delta_0)^{1/2} + C(\delta) = 0$ .
5. Spectrum of the diamagnetic hydrogen atom for unphysical  $D = -2, -4, -6, \dots$ : exact solutions.